

## Guaranteed Content

Water Soluble Calcium Oxide (CaO)

Water Soluble Boron (B)

Chlor Content: %14,66

**%W/W**

%12

%0,3

## DIRECTIONS FOR APPLICATION:

Crop	Application Period	Leaf Application	Soil Application
Fruit Trees	2-3 applications after fruit set	300-500 ml	2-3 lt / decar
Olive	After fruit set at the last 2-3 watering	300-500 ml	2-3 lt / decar
Tomato, Eggplant, Pepper, Cucumber, Zucchini	2-3 applications after fruit set	300-500 ml	2-3 lt / decar
Sugar Beet, Carrot, Potato, Onion	2-3 applications during growth phase	300-500 ml	2-3 lt / decar
Corn, Sunflower, Cotton	Growth phase	300-500 ml	2-3 lt / decar
Grain (Cereal)	After tillering	300-500 ml	2-3 lt / decar
Vineyard	On bunch growth phase and on unripe phase before spotting	300-500 ml	2-3 lt / decar
Citrus	Fruit growth and ripening phases	300-500 ml	2-3 lt / decar
Hazelnut	After fruit set	300-500 ml	2-3 lt / decar
Cotton	2 applications after anchoring	300-500 ml	2-3 lt / decar

## USAGE INSTRUCTIONS:

On all plants; Can be used 2 - 4 lt/da on soil with drip irrigation or sprinkling application. For applications on leaves, 200 ml / 100 L water dosage is advised. Apply rates should be decided considering soil/leaf analyses and plants' necessities. Repeating the application should also be based on analyses and plants' necessities.